

Select	Vaccine Type	Adult Vaccine Recommendation	Dose(s)
<input type="checkbox"/>	Diphtheria provided as Td or Tdap	<ul style="list-style-type: none"> Everyone 	Booster every 10 years
<input type="checkbox"/>	Dukoral®	<ul style="list-style-type: none"> Travellers, depending on destination Adults 18 years of age and over 	<p>Taken orally, 2 doses taken 1-6 weeks apart; last dose taken at least 1 week before travel</p> <p>Provides coverage for traveller's diarrhea for 3 months.</p> <p>Revaccination:</p> <ul style="list-style-type: none"> If up to 5 years since last dose: 1 dose If more than 5 years since last dose: 2 doses
<input type="checkbox"/>	Hepatitis A Avaxim®, Havrix®	<ul style="list-style-type: none"> People with certain medical, occupational, or lifestyle risks Anyone who wants protection from hepatitis A 	<ul style="list-style-type: none"> Avaxim®: initial dose plus 1 follow-up dose 6-36 months later Havrix®: initial dose plus 1 follow-up dose 6 months later
<input type="checkbox"/>	Hepatitis B	<ul style="list-style-type: none"> People with certain medical, occupational, or lifestyle risks Anyone who wants protection from hepatitis B 	<ul style="list-style-type: none"> 3 doses: initial dose plus 2 follow-up doses at 1 and 6 months later <p style="text-align: center;">OR</p> <ul style="list-style-type: none"> 4 doses (rapid schedule): initial dose plus 3 follow-up doses at 7 days, 21 days & 1 year later <p style="text-align: center;">OR</p> <ul style="list-style-type: none"> 4 doses (accelerated schedule): initial dose plus 3 follow-up doses 1, 2 & 12 months later
<input type="checkbox"/>	Herpes zoster (shingles) Shingrix (non-live vaccine), Zostavax® II (live vaccine)	<ul style="list-style-type: none"> Adults 60 years of age and over; may be given to those aged 50 to 59 years 	<ul style="list-style-type: none"> Shingrix: 2 doses (2-6 months apart) Zostavax® II: 1 dose
<input type="checkbox"/>	Human papillomavirus (HPV) (HPV9, Gardasil®9)	<ul style="list-style-type: none"> Women 18 to 45 years of age Men 18 to 26 years of age with lifestyle or other risks Men 27 years of age and older at continued risk of exposure 	3 doses (initial dose plus 2 follow-up doses 2 months and 6 months later)
<input type="checkbox"/>	Influenza (flu)	<ul style="list-style-type: none"> People with chronic medical conditions and those living in close contact with high-risk people All healthy adults, especially those age 65 and over 	Annually
<input type="checkbox"/>	Measles/mumps/rubella (German measles) - MMR	<ul style="list-style-type: none"> Adults born before 1970: 1 dose if travelling, have certain medical conditions, or have occupational risk Adults born in or after 1970 who have not had the disease: 2 doses 	1 or 2 doses
<input type="checkbox"/>	Meningococcal infection ACYW-135 (Menactra®)	<ul style="list-style-type: none"> People with specific medical conditions and those living in communal residences, including students and military personnel 	<ul style="list-style-type: none"> 1 dose Revaccination every 5 years if at continued risk
<input type="checkbox"/>	Meningococcal B (Trumenba®)	<ul style="list-style-type: none"> 18-25 years old People with specific medical conditions and those living in communal residences, including students and military personnel 	<ul style="list-style-type: none"> An initial dose plus a follow-up dose 6 months later; people at high risk: initial dose plus 2 follow-up doses 1 month and 4-6 months later
<input type="checkbox"/>	Pertussis (whooping cough) provided as Tdap	<ul style="list-style-type: none"> Everyone 	Once in adulthood
<input type="checkbox"/>	Pneumococcal infection Pneumococcal Conjugate, (Pevnar®13) – PCV13 Pneumococcal Polysaccharide (Pneumovax®23) – PPSV23	<ul style="list-style-type: none"> Adults 65 years of age and over Immunocompromised adults <ul style="list-style-type: none"> Adults 65 years of age and over Immunocompromised adults People with conditions that put them at high risk (asthma; COPD; diabetes; or chronic lung, kidney, or heart disease) 	<p>Pevnar®13: 1 dose</p> <p>Pneumovax®23: 1 dose plus a booster shot in 5 years for those who are immunocompromised</p>
<input type="checkbox"/>	Tetanus (lockjaw) provided as Td or Tdap	<ul style="list-style-type: none"> Everyone 	Booster every 10 years
<input type="checkbox"/>	Varicella (chickenpox)	<ul style="list-style-type: none"> People who have not had the disease or been vaccinated 	2 doses, at least 4 weeks apart
<input type="checkbox"/>	Additional travel vaccines	<ul style="list-style-type: none"> Vary by destination; consult a travel health clinic or pharmacy with travel health designation, your doctor, or local public health unit 	Vary depending on vaccine